



APPLICATION GUIDELINES

Prosaro® fungicide will help you achieve higher grain quality and more yield so you can claim the crown of higher profit.

- Unsurpassed control of both leaf and head diseases
- Unique combination of fast action and long residual
- No. 1 scab reducer - excellent performance against Fusarium head blight (FHB) and proven deoxynivalenol (DON) reduction in harvested grain
- Proven quality and yield increases under a wide range of environmental conditions

How to Spray

Rate	6.5 – 8.2 fl oz/A
Timing	<ul style="list-style-type: none">• Fusarium Head Blight: Optimal time to apply is as a preventative foliar spray at early flower (Feekes Growth Stage 10.51)• Leaf and Stem Diseases: Apply as a foliar spray when the earliest disease symptoms appear on the leaves and stems
Rainfast	<ul style="list-style-type: none">• 2 hours or until the treated surface is fully dry

Most major leaf and head diseases are well controlled by timely applications of Prosaro.

Spring Wheat / Winter Wheat Diseases Controlled

Fusarium Head Blight (scab)

Leaf, Stem and Stripe Rusts

Powdery Mildew

Septoria Leaf Spot

Glume Blotch

Stagonospora Blotch

Tan Spot

Barley Diseases Controlled

Fusarium Head Blight (scab) – suppression

Net Blotch

Powdery Mildew

Scald

Spot Blotch

Rusts

prosaro.us



BAYER COMMITMENT

Today and tomorrow, Bayer focuses on grower needs and innovation to deliver the most complete and effective corn and crop protection portfolio.

IMPORTANT: This bulletin is not intended to provide adequate information for use of this product. Read the label before using this product. Observe all label directions and precautions while using this product.