



STRATEGO[®] YLD



APPLICATION GUIDELINES

Do all you can to increase your yields and grow your bottom line. Treat your corn and soybeans with **Stratego[®] YLD**. With its advanced technology, **Stratego YLD** delivers:

- All-around plant coverage
- Broad spectrum disease control including strobilurin-resistant strains
- Average yield increase of **7 bu/A** when used early season on corn, **12 bu/A** when used at-tassel on corn and **3-4 bu/A** on soybeans

Corn Diseases*

- Anthracnose leaf blight
- Common rust
- Eyespot
- Gray leaf spot
- Northern corn leaf blight
- Northern corn leaf spot
- Southern corn leaf blight
- Southern rust

Soybean Diseases*

- Alternaria leaf spot
- Anthracnose
- Asian soybean rust
- Cercospora blight
- Frogeye leaf spot
- Pod and stem blight
- Powdery mildew
- Rhizoctonia aerial blight
- Septoria brown spot

*See product label for complete list of diseases controlled.

Product Information

- Chemical Class – Triazole and Strobilurin
- Mode of Action – Inhibition of energy production (QoI) and sterol production (DMI)
- Formulation – SC
- Activity – Preventative and curative
- Residual activity – up to 21 days

APPLICATION INFORMATION

Field Corn (includes Field Corn, Popcorn and Seed Corn)

Use Rate per Application	V4 to V7 – 2 to 2.5 fl oz/A V7 through R2 – 4 to 5 fl oz/A
Use Rate per Season	10 fl oz/A
Timing	V4 to V7 – Apply when conditions are favorable for disease development. V7 through R2 – Apply when disease first appears. A follow-up treatment can be made 7 to 14 days later if conditions for disease development persist. Adjuvants should not be used for applications made between V8 and VT.
PHI	14 days (grain and fodder); 0 days (forage)

Sweet Corn

Use Rate per Application	4 to 5 fl oz/A
Use Rate per Season	20 fl oz/A
Timing	Apply when disease first appears and continue on a 5- to 14-day interval if favorable conditions for disease development persist.
PHI	0 days (forage and ear); 14 days (fodder)

Soybeans

Use Rate per Application	4 to 4.65 fl oz/A
Use Rate per Season	13.95 fl oz/A
Timing	Apply at early flowering or prior to disease development, whichever is earlier. Repeat application on a 10- to 21-day spray interval if conditions are favorable for continued disease development.
PHI	21 days

IMPORTANT: This bulletin is not intended to provide adequate information for use of this product. Read the label before using this product. Observe all label directions and precautions while using this product.

