























PEANUT RECOMMENDED APPLICATION TIMING

PESTS	PRODUCT	AT PLANTING	30 DAYS	45 DAYS	60 DAYS	75 DAYS	90 DAYS	105 DAYS	120 DAYS
APHIDS, LEAFHOPPERS, THRIPS, WHITEFLIES									
									
NEMATODES (SUPPRESSION)									
									
CYLINDROCLADIUM BLACK ROT (SUPPRESSION), RHIZOCTONIA, WHITE MOLD									
									
									
LEAF SPOT									
									
									
									

*In the case of Velum Total, suppression only.

 Insecticides/Nematicides  Fungicides

IMPORTANT: This bulletin is not intended to provide adequate information for use of these products. Read the label before using these products. Observe all label directions and precautions while using these products.







For more information, visit www.CropScience.Bayer.us.

ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. Not all products are registered in all states and may be subject to use restrictions. The distribution, sale, or use of an unregistered pesticide is a violation of federal and/or state law and is strictly prohibited. Check with your local dealer or representative for the product registration status in your state. Absolute®, Admire®, Bayer, Bayer Cross, Proline®, Propulse®, Provost® and Velum® are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2021 Bayer Group. All rights reserved. 1T022110097



The Bayer Peanut Protection Portfolio

Bayer provides solutions to help you grow high-quality peanuts, increase yields and continuously improve the vigor of your peanut crop.

	PRODUCT	KEY PESTS/DISEASES CONTROLLED	CHEMICAL CLASS GROUP NO.	REI (HOURS)	PHI (DAYS)	USE RATE*
INSECTICIDES/NEMATOCIDES	 <p>Soil Application A systemic insecticide that provides economical and enduring control of damaging insects.</p>	Aphids, Leafhoppers, Thrips (suppression), Whiteflies	Neonicotinoids IRAC Group 4A	5 days	14	7 to 10.5 fl. oz./A
	<p>Foliar Application A systemic insecticide that provides economical and enduring control of damaging insects.</p>	Aphids, Leafhoppers, Thrips (suppression), Whiteflies	Neonicotinoids IRAC Group 4A	5 days	14	1.2 fl. oz./A
	 <p>Moves from the plant's roots to the leaves, not only suppressing nematodes below ground, but also moving throughout the plant to help protect root health, suppress key diseases and maximize yield potential.</p>	Aphids, Early leaf spot (suppression), Late leaf spot (suppression), Leafhoppers, Nematodes (suppression), Thrips, Whiteflies, White mold (suppression)	FRAC Group 7	12	7	6.5 to 6.84 fl. oz./A
FUNGICIDES	 <p>Effectively and efficiently controls major fungal diseases while also promoting all-around plant health to maximize yield potential.</p>	Early leaf spot, Late leaf spot, Rhizoctonia, Rust, Web blotch	FRAC Group 11 (QoI) FRAC Group 3 (DMI)	12	14	3.5 to 7 fl. oz./A
	 <p>Provides powerful, broad-spectrum disease protection to maximize yield potential.</p>	Cylindrocladium black rot (suppression), Early leaf spot, Late leaf spot, Rhizoctonia, Southern blight, White mold	FRAC Group 3 (DMI)	12	14	5.7 fl. oz./A (in-furrow and banded)
	 <p>Protects against key diseases throughout the growing season, resulting in improved plant health as well as high-quality crops, and provides an effective approach to resistance management.</p>	Cylindrocladium black rot (suppression), Early leaf spot, Late leaf spot, Nematodes, Rhizoctonia, Rust, Sclerotinia blight, Southern blight, White mold	FRAC Group 7 (SDHI) FRAC Group 3 (DMI)	12	14	13.6 fl. oz./A
	 <p>Combines excellent protection against key soil and foliar diseases with the convenience of tankmix flexibility.</p>	Cylindrocladium black rot (suppression), Early leaf spot, Late leaf spot, Leaf rust, Leaf scorch and pepper spot, Rhizoctonia, Southern blight, Web blotch, White mold	FRAC Group 3 (DMI)	12	14	11 to 13 fl. oz./A

*See product label for specific rates for target pests.