
POTATO RECOMMENDED APPLICATION TIMING



IMPORTANT: This bulletin is not intended to provide adequate information for use of these products. Read the label before using these products. Observe all label directions and precautions while using these products.

For more information, visit www.CropScience.Bayer.us

Signal word for Baythroid® XL and Leverage® 360: WARNING. Signal word for all other products: CAUTION.





The Bayer Potato Protection Portfolio

Bayer provides solutions to help you grow high-quality potatoes, increase yields and continuously improve the vigor and marketability of your potato crop.

		PRODUCT	KEY PESTS/DISEASES CONTROLLED	CHEMICAL CLASS GROUP NO.	REI (HOURS)	PHI (DAYS)	USE RATE*
INSECTICIDES/MITICIDES/NEMATICIDES	ADMIRE° PRO	A soil-applied systemic insecticide that provides economical and enduring control of damaging insects.	Aphids, Colorado potato beetles, Flea beetles, Psyllids, Wireworms (seed piece treatment)	Neonicotinoids IRAC Group 4A	12	N/A	Soil: 5.7 to 8.7 fl. oz./A Seed piece treatment: 3.5 to 7 fl. oz./A
	BAYTHROID XL	Quickly knocks down a wide range of pests, provides long-lasting residual and high return on investment.	Cabbage loopers, Colorado potato beetles, Cutworms	Pyrethroids IRAC Group 3	12	0	0.8 to 2.8 fl. oz./A
	LEVERAGE 360 †	Features two modes of action against harmful sucking and chewing pests – one delivers fast knockdown, the other extends residual control.	Aphids, Cabbage loopers, Colorado potato beetles	Neonicotinoids IRAC Group 4A Pyrethroids IRAC Group 3	12	7	2.8 fl. oz./A [†]
	MOVENTO HL	Features powerful, two-way movement throughout the plant to protect it from a broad range of insects, mites and nematodes.	Aphids, Mites (suppression), Nematodes (suppression), Psyllids, Wireworms (suppression)	Tetramic acids IRAC Group 23	24	7	2 to 2.5 fl. oz./A
TICIDES	o·b·e·r·o·n	Provides excellent control of mites and psyllids but soft on beneficials, making it a good miticide choice that fits well in IPM systems.	Psyllids, Twospotted spider mites	Tetronic acids IRAC Group 23	12	7	4 to 8 fl. oz./A (Oberon® 4 SC)
INSECT	SIVANTO° prime	Precisely targets key damaging pests while helping safeguard beneficial insects and mites, including honey bees, to preserve the overall health of your plants and your investment.	Aphids, Colorado potato beetles, Leafhoppers, Potato psyllids, Whiteflies	Butenolides IRAC Group 4D	4 (12 in CA)	7	7 to 14 fl. oz./A
	VELUM ° PRIME	Moves from the plant's roots to the leaves, not only suppressing nematodes below ground, but also moving throughout the plant to help protect root health, suppress key diseases and maximize yield potential.	Nematodes (suppression), Early blight (suppression), White mold (suppression)	FRAC Group 7	12	7	6.5 to 6.84 fl. oz./A
ES	Luna	Protects against key diseases throughout the growing season, resulting in improved plant health as well as high-quality crops, and provides an effective approach to resistance management.	Black dot, Botrytis, Brown spot, Early blight, White mold	FRAC Group 7 (SDHI) FRAC Group 9 (AP)	12	7	8 to 11.2 fl. oz./A
FUNGICIDES	Previcur	Provides proven control of damaging diseases by quickly penetrating the leaf surface and moving throughout the plant to protect new growth.	Late blight	Carbamates FRAC Group 28	12	14	0.7 to 1.2 pt./A + tankmix partner
-	SCALA	Provides high level of control against many economically damaging diseases and helps control many resistant disease strains.	Botrytis, Brown spot, Early blight	FRAC Group 9 (AP)	12	7	7 fl. oz./A
	MINUET	Minuet is a soil-applied biological. In the soil, it forms a symbiosis with the plant and triggers activation of root and plant growth as well as the plant defense response to support healthy plants and increased crop quality and yield potential.	Pythium, Rhizoctonia, Verticillium	Bacillus subtilis, QST 713 FRAC Group 44	4	0	12 to 24 fl.oz./A
BIOLOGICALS	SERENADE ASO	Provides protection against soil and foliar fungal and bacterial diseases through multiple sites of action with a flexible application.	Soil Application: Erwinia spp., Fusarium spp., Phytophthora spp., Pythium spp., Rhizoctonia spp., Verticillium spp.	Bacillus subtilis QST 713 FRAC Group 44	4	0	2 to 4 qt./A
			Foliar Application: Black dot, Botrytis, Early Blight, White Mold				
	SERENADE	Provides protection against foliar fungal and bacterial diseases through multiple sites of action with a flexible application.	Black rot, Botrytis, White Mold (suppression)	Bacillus subtilis QST 713 FRAC Group 44	4	0	14 to 20 oz./A
SEED PIECE TREATMENT	EMESTO°	An innovative seed treatment fungicide that helps manage seed and soil-borne disease and promotes early crop establishment and seed vigor.	Fusarium tuber rot (suppression), Seed and soil-borne Rhizoctonia (suppression), Silver scurf (suppression)	FRAC Group 7 (SDHI) FRAC Group 3 (DMI)	12	N/A	0.3 fl. oz./cwt

^{*}See product label for specific rates for target pests. **Baythroid® XL and Leverage® 360 are Resticted Use Pesticides. †Do not use if Admire® Pro was used at planting. Do not use at bloom.